

Erratum to: A multidisciplinary approach to enhance the conservation and use of hazelnut *Corylus avellana* L. genetic resources

L. Bacchetta · M. Rovira · C. Tronci · M. Aramini ·
P. Drogoudi · A. P. Silva · A. Solar · D. Avanzato ·
R. Botta · N. Valentini · P. Boccacci

Received: 1 January 2014 / Accepted: 27 August 2014 / Published online: 9 May 2015
© Springer Science+Business Media Dordrecht 2015

Erratum to: Genet Resour Crop Evol
DOI 10.1007/s10722-014-0173-7

We regret that the table part of Fig. 3 was missed in the online publication. The correct figure is given below.

The online version of the original article can be found under doi:[10.1007/s10722-014-0173-7](https://doi.org/10.1007/s10722-014-0173-7).

L. Bacchetta (✉) · C. Tronci · M. Aramini
Agenzia Nazionale per le Nuove Tecnologie, l'Energia e lo Sviluppo Economico Sostenibile, ENEA Casaccia - UTAGRI, Via Anguillarese, 301, 000123 Rome, Italy
e-mail: loretta.bacchetta@enea.it

M. Rovira
Division IRTA, Institut de Recerca i Tecnologia Agroalimentàries, Centre Mas de Bover, Ctra. Reus-El Morell, km 3.8, 43120 Constantí, Tarragona, Spain

P. Drogoudi
Pomology Institute, Hellenic Agricultural Organization 'Demeter', D.G. of Agricultural Research, 38 R.R. Station, 59 035 Naoussa, Greece


A. P. Silva
CITAB, Centre for the Research and Technology of Agro-Environment and Biological Sciences, Universidade de Trás-os-Montes e Alto Douro, Apartado 1013, 5001-801 Vila Real, Portugal

A. Solar
Biotehniska Fakulteta, Univerza v Ljubljani, Jamnikarjeva 101, 1000 Ljubljana, Slovenia

D. Avanzato
CRA Centro Ricerca per la Frutticoltura, Via Fioranello, 52, 0013 Rome, Italy

R. Botta · N. Valentini
Department of Agricultural, Forestry and Food Sciences (DISAFA), University of Turin, Largo Paolo Braccini, 2, 10095 Grugliasco, TO, Italy

P. Boccacci
Institute for Sustainable Plant Protection - National Research Council (IPSP-CNR), UOS of Grugliasco, Largo Paolo Braccini, 2, 10095 Grugliasco, TO, Italy



Variable	PC1	PC2	PC3
oil content	-0.514	-0.329	-0.259
%			
C16:0	-0.302	0.833	-0.102
C16:1	-0.16	0.597	-0.433
C18:0	-0.68	0.381	0.24
C18:1	-0.737	-0.649	0.068
C18:2	0.918	0.343	-0.095
C18:3	0.485	0.178	0.185
SFA	-0.585	0.800	0.058
Total	-0.742	-0.642	0.062
MUFA			
Total PUFA	0.92	0.343	-0.09
UFA/SFA	0.625	-0.765	-0.065
MUFA/SFA	0.362	-0.913	-0.03
PUFA/SFA	0.98	0.041	-0.125
alfa-	0.258	-0.084	0.91
tocopherol			
SI	0.087	0.259	0.922
Eigenvalue	5.759	4.538	2.085
Proportion	38.395	30.257	13.897
of total (%)			

Fig. 3 Position of the principal component (PC) scores of the hazelnut kernel composition for 75 hazelnut cultivars. Numbers refer to the clone number of the cultivar: 1 *C.a.* 9 Raul; 2 Gunslebert; 3 Barcellona; 4 Hall's Giant; 5 Segorbe; 6 Dal Rosso; 7 *C. maxima* à Pellicule Blanche; 8 Negret; 9 Fertile de Coutard; 10 Bergeri; 11 CV/2; 12 Rotblattrige Lambertnuss; 13 *C. maxima* à Pellicule Rose; 14 Merveille de Bollwiller; 15 Longue d'Espagne; 16 Pautet (ref. cv.); 17 Incrocio L35; 18 Trenet; 19 Vermellet; 20 Corabel; 21 Casina; 22 Nostrale; 23 Provence; 24 Pallagrossa; 25 CV/1; 26 Tonda di Giffoni (ref. cv.); 27 Pellicola bianca; 28 San Giovanni; 29 Tonda bianca; 30 Ferwiller; 31 Istrska okrogloplodna leska; 32 Tonda Gentile Langhe (ref. cv.);

33 Feriale; 34 Polycarpus wild; 35 San Vicino; 36 Molar; 37 Lunga Ginnasi; 38 Gironell; 39 Tonda di Biglini; 40 TGL d(clone PD); 41 *C.a.* 5 Grada de Visu; 42 Cosford; 43 *C.a.* 11 Tubulosa; 44 Istrska dolgoplodna leska; 45 Nocchione; 46 Imperatrice Eugenie; 47 *C.a.* 12 Purpurea; 48 *C.a.* 7 DaVeiga; 49 GR pi 03; 50 Piazza armerina; 51 Nociara; 52 Riccia di Talanico; 53 Meloni; 54 Karydato; 55 Carrello; 56 Tonda gentile romana; 57 *Avellana* Speciale; 58 Argiroupoli; 59 Patem; 60 Palaz; 61 Ada; 62 Barettona Vico; 63 *C.a.* 8 Comum; 64 Castanyera (ref. cv.); 65 Extra Giaghli; 66 Morell; 67 Grifoll; 68 Camponica; 69 Daria; Tombul Giaghli; 71 Comune di Sicilia; 72 Tonda Calabrese-Caserta; 73 Barettona; 74 Centenaria Ginnasi; 75 Culpia